



Microparticles and their impact on intestinal immunity

Becker, Helen M ; Bertschinger, Martina M ; Rogler, Gerhard

Abstract: Microparticles are small ($<1\ \mu\text{m}$), nonbiological particles that are used in many areas of daily life. As food additive they are used as anticaking agents or food colorants. The most common food-derived ingested compounds are aluminium silicate and titanium dioxide (TiO(2)), the latter being a white pigment used in toothpaste or sugar toppings. The increasing abundance of microparticles in the Western diet raises the question of the potential risks associated with gastrointestinal diseases such as Crohn's disease (CD). Accumulation of particles has been shown in cells of Peyer's patches, but it is not clear whether this also has pathological effects. NLRP3 is a member of the intracellular pattern recognition receptor family and it is part of the inflammasome, a multiprotein complex containing caspase-1 which activates the proinflammatory cytokines interleukin (IL)-1 and IL-18. With regard to recent findings identifying small particles such as asbestos and monosodium urate as NLRP3 activators, TiO(2) may be another potential target for inflammasome studies. We found that macrophage-like cells readily take up TiO(2) after 6 h. Incubation of cells with TiO(2) resulted in the assembly of NLRP3 with caspase-1. This inflammasome assembly correlated with secretion of IL-1. In intestinal epithelial cells, TiO(2) also was found to be ingested. The counting of particles localized intracellularly revealed a dose-dependent increase of TiO(2)-positive cells. This points to the fact that in humans with a leaky intestinal barrier (such as IBD patients), TiO(2) microparticles may be taken up by macrophages and intestinal epithelial cells, may activate the inflammasome and induce IL-1 and IL-18 secretion. This may aggravate inflammation in susceptible individuals.

DOI: <https://doi.org/10.1159/000342602>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-76197>

Journal Article

Published Version

Originally published at:

Becker, Helen M; Bertschinger, Martina M; Rogler, Gerhard (2012). Microparticles and their impact on intestinal immunity. *Digestive Diseases*, (30 Suppl.):47-54.

DOI: <https://doi.org/10.1159/000342602>

Microparticles and Their Impact on Intestinal Immunity

Helen M. Becker^a Martina M. Bertschinger^b Gerhard Rogler^b^aDepartment of Medicine, University of Calgary, Calgary, Alta., Canada; ^bDivision of Gastroenterology and Hepatology, University Hospital of Zurich, Zurich, Switzerland

Key Words

Inflammasome · Microparticles · Interleukin-18

Abstract

Microparticles are small (<1 µm), nonbiological particles that are used in many areas of daily life. As food additive they are used as anticaking agents or food colorants. The most common food-derived ingested compounds are aluminium silicate and titanium dioxide (TiO₂), the latter being a white pigment used in toothpaste or sugar toppings. The increasing abundance of microparticles in the Western diet raises the question of the potential risks associated with gastrointestinal diseases such as Crohn's disease (CD). Accumulation of particles has been shown in cells of Peyer's patches, but it is not clear whether this also has pathological effects. NLRP3 is a member of the intracellular pattern recognition receptor family and it is part of the inflammasome, a multiprotein complex containing caspase-1 which activates the proinflammatory cytokines interleukin (IL)-1β and IL-18. With regard to recent findings identifying small particles such as asbestos and monosodium urate as NLRP3 activators, TiO₂ may be another potential target for inflammasome studies. We found that macrophage-like cells readily take up TiO₂ after 6 h. Incubation of cells with TiO₂ resulted in the assembly of NLRP3 with caspase-1. This inflammasome assembly correlated with secretion of IL-1β. In intestinal epithelial cells, TiO₂ also was found to be ingested. The counting of particles lo-

calized intracellularly revealed a dose-dependent increase of TiO₂-positive cells. This points to the fact that in humans with a leaky intestinal barrier (such as IBD patients), TiO₂ microparticles may be taken up by macrophages and intestinal epithelial cells, may activate the inflammasome and induce IL-1β and IL-18 secretion. This may aggravate inflammation in susceptible individuals.

Copyright © 2012 S. Karger AG, Basel

Introduction

Intestinal epithelial cells (IECs) form a single layer on the surface of the intestinal mucosa and the primary barrier against the content of the gut lumen. They protect the underlying tissue and the entire organism from potentially harmful microorganisms, particles or larger molecules. IECs secrete antimicrobial substances such as defensins [1, 2] and communicate with the immune cells of the intestinal mucosa via a high number of soluble mediators such as chemokines and cytokines. In patients with inflammatory bowel disease (IBD), the epithelial and mucosal barrier function is disturbed, finally leading to a chronic relapsing inflammation of the intestinal mucosa.

Both environmental triggers and genetic susceptibility have been shown to contribute to the pathophysiology of IBD. More than 140 susceptibility loci for IBD have been identified mainly by genome-wide association stud-

ies (GWAS) [3], among them a number of genes related to innate immune functions including single-nucleotide polymorphisms (SNPs) in so-called pattern-recognition receptors (PRRs) which recognize microbial patterns (i.e. pathogen-associated molecular patterns, PAMPs) but also other molecular signatures that may be a risk for health (so-called danger-associated molecular patterns, DAMPs).

In healthy individuals, activation of PRRs and the subsequent proinflammatory signalling is tightly regulated. Among other members of the PRR family, NLRP3 [nucleotide-binding domain-containing and leucine-rich repeat-containing (NLR) family and pyrin domain-containing 3 (PYD)], has been shown to play a role in IBD etiology. Alternative names for NLRP3 are NALP3, CIAS (cold-induced autoinflammatory syndrome 1), PYPAF (PYD-containing Apaf1-like) and cryopyrin. In biopsies of Crohn's disease (CD) patients, NLRP3 expression was found to be increased [4]. Besides microbial patterns, NLRP3 also recognizes 'danger signals' or DAMPs such as adenosine triphosphate (ATP) [5], asbestos [6] and uric acid crystals [7]. Upon activation signals, NLRP3 gets associated with a large multiprotein complex, the NLRP3 inflammasome, which activates caspase-1. Activated caspase-1 then finally cleaves pro-interleukin (IL)-1 β and pro-IL-18 (fig. 1).

Expression of all components of the NLRP3 inflammasome has been observed in professional immune cells such as macrophages or dendritic cells, but a high level has also been found in IECs.

NLRP3 and Susceptibility to IBD

Similar to nucleotide-binding oligomerization domain-containing protein 2 (NOD2), the first-identified CD susceptibility gene, NLRP3, contains a 'caspase-recruitment domain' (CARD) and, upon activation, assembles with caspase-1 to form the inflammasome [8]. The relevance of NLRP3 for CD pathogenesis is obvious from the finding that a gain-of-function SNP in NLRP3 and a correlating CARD8 loss-of-function SNP both result in a protective effect against CD [9, 10]. NLRP3 gain-of-function mutations were previously associated with other inflammatory diseases and syndromes such as rheumatoid arthritis [11], Muckle-Wells syndrome, familial cold autoinflammatory syndrome [12] and chronic infantile neurological cutaneous and articular syndrome [13]. All are characterized by elevated IL-1 β levels, indicating an NLRP3/inflammasome activation. Further, NLRP3 was




Fig. 1. NLRP3 inflammasome activation. PAMPs and DAMPs such as ROS have been shown to activate the NLRP3 inflammasome. Upon activation, NLRP3 assembles to a multiprotein complex together with the adaptor protein, apoptosis-like speck protein (Asc) and caspase-1. Caspase-1 undergoes autocatalytic activation, which enables the cleavage of its substrates pro-IL-1 β and pro-IL-18. The cleaved cytokines are then released by the cell to activate downstream targets.

shown to be upregulated in colon tissue after the induction of acute or chronic colitis in mouse models. An elevated expression was also found in colon specimens of CD patients [4].

The inflammasome proteins belong to a family of NLRs, all sharing common domains that act as intracellular sensors for PAMPs and DAMPs. Similar to Toll-like receptors, the NLR family is evolutionarily ancient and shares structural homologies with plant resistance (R-) proteins. Both protein families have leucine-rich repeat (LRR) regions for the recognition of bacterial motifs, an oligomerization domain and an effector domain [14] (fig. 2).

Upon activation, NLRP3 assembles into a multiprotein complex, recruiting other NLRP3 proteins, CARD8, ASC (apoptosis-like speck protein) and caspase-1 (fig. 1). The recruitment of caspase-1 triggers its autocatalytic cleavage and activation. Subsequently, caspase-1 cleaves pro-IL-1 β or pro-IL-18; this is usually followed by the release of biologically active forms of these cytokines (table 1).




Fig. 2. Domains of the NLR family. In the NLRP subfamily, the PYD enables interaction with other proteins, whereas NOD1, NOD2, ice protease-activating factor (IPAF) and MHC class II transactivator (CIITA) contain CARD domains for interaction. Furthermore, LRRs are found at the C-terminus of all NLR family members. AD = Acid transactivation domain; BIR = baculovirus inhibitor of apoptosis repeat; FIIND = domain with function to find; NACHT = domain present in NAIP, CIITA, HET-E, and TP-1; NAD = NBD-associated domain; NAIP = NLR family, apoptosis inhibitory protein; NALP = NLRP [66].

Food-Derived Environmental Factors in the Pathogenesis of IBD?

IBD cannot only be attributed to genetic factors, as monozygotic twins have a concordance rate of only 20–50% of developing IBD [34–37]. The increased incidence of IBD in the last 100 years, the higher incidence of IBD in countries with a Western lifestyle and the observed rise of IBD in countries recently undergoing ‘westernization’ in Asia have led to several hypotheses explaining this phenomenon, summarized by Hanauer [38]. Hygiene, diet, social status and smoking are factors potentially influencing the risk for developing IBD [39]. Some childhood factors such as breastfeeding and childhood infections are likely to have protective functions, and it has been suggested that increased hygienic standards increase the risk of IBD. Another factor associated with a typical Western diet is microparticles.

Table 1. Activators of the NLRP3 inflammasome (table adapted from [15])

		Reference
<i>Microorganisms</i>		
Sendai virus	09/2006	[16]
Influenza virus	09/2006, 04/2009	[16, 17]
Adenovirus	03/2008	[18]
<i>Staphylococcus aureus</i>	03/2006, 09/2009	[5, 19]
<i>Listeria monocytogenes</i>	03/2006, 05/2008	[5, 20]
<i>Saccharomyces cerevisiae</i>	06/2009	[21]
<i>Candida albicans</i>	08/2009	[22]
<i>Aspergillus fumigatus</i>	04/2010	[23]
Bacterial RNA or synthetic purine bases	03/2006	[24]
Bacterial pore-forming toxins (aerolysin, maitotoxin and nigericin)	03/2006, 09/2006, 11/2006	[5, 25, 26]
Malaria toxin	08/2009	[27]
<i>Danger signals</i>		
Extracellular ATP	03/2006	[5]
Hyaluronan	03/2009	[28]
Glucose	12/2009	[29]
Monosodium urate	03/2006	[7]
Amyloid-beta	08/2008	[30]
Silica	08/2008, 07/2008	[31, 32]
Asbestos	07/2008, 05/2008	[6, 32]
Alum	08/2008	[31]
Cholesterol crystals	04/2010	[33]

Microparticles are used as food additives in order to influence the color, the consistency or the appearance of a product. The most commonly used additives are titanium dioxide (TiO₂, E171) and aluminium silicate (AlSi, E559). Both compounds are sub-micron-sized (0.1–1 µm diameter) oxides of titanium, silicon or aluminium. Since they are inorganic, they are highly stable and resistant to degradation. According to current European legislation, there is no maximum TiO₂ level specified. It is recommended that TiO₂ should be used in amounts according to good manufacturing practice [40]. The international agency for research on cancer published an overview on studies addressing the hazardous potential of TiO₂ in 2006. In animal studies, inhalation was followed by the development of tumors. In vitro studies with human alveolar macrophages revealed inhibited phagocytosis and DNA damage. Subsequently, TiO₂ was rated as possibly carcinogenic for humans (Group 2b) [41]. Classification by the International Agency for the

Research on Cancer, IARC, as Group 2B. This means that the agent is possible carcinogenic in humans but there is limited evidence from either animal experiments or human studies. In the case of TiO₂ there is evidence from animal studies but limited evidence from human studies.

Different Microparticles Used in the Food Industry

TiO₂ is a white, crystalline powder which is widely used as a food supplement in refined food. The pigment is used in toothpaste, confectionery, white sauces and dressings and nondairy creamers [42]. In the pharmaceutical industry, it is used as a pill coating. It is inert to most influences (fig. 3).

Iron Oxide

Iron oxide (Fe₂O₃) is a nonsoluble, brown and red pigment. It is used as a food colorant as well as in cosmetics. The product name is 'Red luster' which is also used for coating pharmaceutical formulations.

Titanium Dioxide/Iron Oxide Mixture

TiO₂/Fe₂O₃ mixture is a gold mica powder. The powder is used as a food colorant and as a coating for pharmaceutical formulations.

AlSi

AlSi, also known as kaolin, is a natural white, fine-grained material. The field of application includes the paper-manufacturing industry and powder fabrication, but it is also used in the food industry as a carrier substance, bleaching product and parting agent, as it prevents caking (fig. 3).

Titanium Oxide/AlSi Mixture

TiO₂ with potassium (K⁺) and AlSi added is used as a whitener (fig. 3).

Sources of Dietary Microparticles

Microparticles are not declared in the contents of many products. The daily intake is difficult to evaluate and is highly dependent on the individual diet. According to a study by the European Food Safety Authorities (EFSA), the estimated daily intake of TiO₂ is 70–80 mg/day [43]. Sources may be tablets, milk whitener and toothpaste among others. Other authors estimate the daily ingested amount of both TiO₂ and AlSi to be 40 mg or 10¹² particles/person/day [44, 45].

An overview of sources and estimated amounts of microparticles is given in table 2 adapted from [43, 44].

Effects of Titanium Dioxide on Intestinal Epithelial Cells

Microparticles may penetrate the intestinal barrier and accumulate in the mucosa. Early reports on dark, pigmented structures in human intestinal lymphoid aggregates date back to 1987 [46]. These structures were later found to contain high amounts of titanium [47]. TiO₂ particles have been shown to accumulate in M cells of Peyer's patches and are passed on to underlying macrophages [48]. Interestingly, the sites of TiO₂ uptake correspond to regions where the first signs of inflammation in CD manifest [49].

In vivo studies on the capacity of TiO₂ to penetrate the gastrointestinal tract revealed that TiO₂ can be found in systemic organs after an oral exposure of 10 days [50–52]. In humans, after a single ingestion of 22 mg and 45 mg of TiO₂, Ti levels in the blood rose after 0.5–2 h and reached a maximum after 8–12 h [52].

Clinical studies have also addressed the influence on TiO₂ on CD patients. A pilot study performed in 2002 observed significant decreases in the CD activity index [53] after 4 months on a low-TiO₂ diet. However, a larger multicenter double-blind trial was not able to confirm these results [54]. Under present living conditions, a reduction or exclusion of dietary TiO₂ may be very difficult. With the recent discovery of it being an activator of the NLRP3 inflammasome, the influence of TiO₂ on IEC in the healthy and in the chronically inflamed mucosa has gained new interest. Dendritic cells derived from NLRP3-deficient mice were shown to secrete less IL-1β upon exposure to TiO₂ [55]. In human keratinocytes and in murine macrophages, TiO₂ activated the NLRP3 inflammasome [56]. These data point towards a role of the inflammasome as a direct or indirect sensor of TiO₂.




Fig. 3. Different microparticles in light microscopy. The examined microparticles TiO_2 (a), a mixture of TiO_2 and iron oxide (b), iron oxide (c), AlSi (d) and a mixture of TiO_2 and AlSi, were diluted to 100 $\mu\text{g}/\text{ml}$ and investigated with a light microscope. Scale bar: 100 μm .

Pathways of NLRP3 Activation

It is still not clear how the NLRP3 inflammasome is activated in detail. Some authors suggest an initial priming event via the NF- κB pathway [57]. The so-called DAMPs comprise a group of inflammasome activators, and several pathways and mechanisms have been shown to be involved in a multistep activation. Recently, Chen and Nunez [58] summarized the most important facts on DAMP-associated inflammasome activation. Three separate pathways have been suggested. The first activation pathway requires high levels of extracellular ATP. ATP may be released by apoptotic/necrotic cells and trigger the opening of the purinergic receptor P2X, ligand-gated ion channel 7 (P2RX7). This channel triggers the formation of pores in the cell membrane that allow an efflux of K^+ ions. Inflammasome activation can be prevented by high concentrations of extracellular K^+ [32, 59].

Table 2. Sources of microparticles in food products and estimated daily intake

Source	mg/person/day
TiO_2	
Food supplement tablet	37.5
Confectionary	24.4
Medicinal product tablet	15.0
Coffee whitener	0.52
Hard coated candies	0.32
Chewing gum	0.28
Marshmallows	0.27
Low-fat or fat-free dressings	0.22
AlSi	
Salt	1.30
Drinking chocolate powder	1.26
Chewing gum	0.92
Sugar, icing	0.30




Fig. 4. Eosin-stained cells stimulated with TiO_2 . TiO_2 particles could be found intracellularly. The most frequent uptake of particles could be detected in cells at the margin of the monolayer. Scale bars: 50 μm (a, b) and 20 μm (c).

The second pathway may be relevant for uric acid crystals (similar to cholesterol crystals), which induce inflammasome activation by a different mechanism [7, 33]. These crystals are taken in lysosomes by cells such as macrophages. As they cannot be degraded, lysosomal damage occurs followed by release of the lysosomal protease cathepsin B which, in turn, triggers NLRP3 activation. The lysosomal rupture may cause the production of reactive oxygen species (ROS) [60, 61]. In line with this, the activation of the inflammasome, and subsequently caspase-1, in response to asbestos crystals was partially prevented when ROS synthesis was inhibited [6]. Release of ROS is a danger signal for surrounding cells, and NLRP3 has been shown to be activated by necrotic cells. In the third concept of inflammasome activation, ROS is regarded to be the most important trigger. ROS may act both as extracellular and intracellular danger signals.

Microparticles and Inflammasome Activation

Under *in vitro* conditions we found microparticles consisting of TiO_2 intracellularly in IEC and macrophages already at low concentrations (fig. 4). Their intracellular accumulation did not depend on endocytosis. TiO_2 aggregates were not restricted to a specific subcellular structure as observed by electron microscopy. The incubation of macrophages and IEC with low concentrations of TiO_2 resulted in assembly of the NLRP3 inflammasome, and correspondingly, the downstream effectors caspase-1, IL-1 β and IL-18 were activated and secreted. High doses of

TiO_2 caused the generation of ROS in the tested cell types, as previously reported for other particulate structures [6]. In contrast to other findings with silica crystals [31], the uptake of TiO_2 did not lead to lysosomal rupture.

Increased levels of functionally active IL-18 in IEC from CD patients are indicative of an increased inflammasome activation. This is in line with findings of other groups reporting the transcriptional upregulation of IL-18 and increased protein levels of mature IL-18 in the mucosa of CD patients [62, 63].

TiO_2 has thus far mainly been the subject of studies focusing on effects of inhalation exposure. However, microparticles are also ingested on a daily basis in the Western diet and their use as food additive is on the rise and not limited by food laws. The long-term impact of dietary microparticles such as TiO_2 and AlSi has not been studied, and the risk evaluation is inconclusive. Given that TiO_2 is also used in pharmaceutical pills, one can assume that local luminal concentrations of TiO_2 can reach substantial amounts. Depending on the properties of the pills, they disintegrate in different sections of the gastrointestinal tract and may affect the intestinal epithelium in a very distinct and concentrated manner.

Further work should include other dietary microparticles such as AlSi and Fe_2O_3 and evaluate their capacity to activate the NLRP3 inflammasome and proinflammatory signalling in macrophages and in IEC. Preliminary data show that, for instance, Fe_2O_3 may be too large to penetrate the cells. An effect of TiO_2 on epithelial cells that has been reported by others was the reduction of transepithelial resistance [64, 65].

Excluding microparticles from food could be a way to improve the well-being of IBD patients. Therefore, it would be worthwhile to initiate further studies with IBD patients to reevaluate the impact of dietary microparticles on the disease activity.

Conclusion

The immune function of IEC is a large field of research, and it has become evident that these cells play an important role as a first line of defense, but also function as signal inducers for underlying immune cells. The tight regulation of proinflammatory and anti-inflammatory signalling is an important feature of IEC, which is out of balance in the state of chronic intestinal inflammations. The emerging role of genetic variations leading to an increased susceptibility to develop IBD is overwhelming, but the actions of many susceptibility genes are not fully understood.

The discovery of NLRP3 as a platform for the recognition of various microorganisms and the more recent revelations on the nonbacterial activators of NLRP3 have gained much attention in recent years. The identification of TiO₂ microparticles as activators of the NLRP3 inflammasome shows that not only immune cells, but also IEC are responsive to nonbacterial stimuli. The current data suggest that TiO₂ triggers inflammation, ROS and, at high concentrations, also apoptosis. Since TiO₂ is consumed as a white pigment in a number of foods and pharmaceutical pills, this finding may evoke questions about food safety, especially in the context of IBD. Further investigation in animal models but also in healthy individuals and IBD patients will help to reevaluate the risk of microparticles in our food.

Disclosure Statement

G.R. discloses grant support from Abbot, Ardeypharm, Essex, FALK, Flamentera, Novartis, Tillots, UCB, Vifor and Zeller.

References

- Jones DE, Bevins CL: Paneth cells of the human small intestine express an antimicrobial peptide gene. *J Biol Chem* 1992;267:23216–23225.
- O'Neil DA, Porter EM, Elewaut D, et al: Expression and regulation of the human beta-defensins hBD-1 and hBD-2 in intestinal epithelium. *J Immunol* 1999;163:6718–6724.
- Franke A, McGovern DP, Barrett JC, et al: Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's disease susceptibility loci. *Nat Genet* 2010;42:1118–1125.
- Villani AC, Lemire M, Fortin G, et al: Common variants in the NLRP3 region contribute to Crohn's disease susceptibility. *Nat Genet* 2009;41:71–76.
- Mariathasan S, Weiss DS, Newton K, et al: Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 2006;440:228–232.
- Dostert C, Petrilli V, Van Bruggen R, Steele C, Mossman BT, Tschopp J: Innate immune activation through NALP3 inflammasome sensing of asbestos and silica. *Science* 2008;320:674–677.
- Martinon F, Petrilli V, Mayor A, Tardivel A, Tschopp J: Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 2006;440:237–241.
- Agostini L, Martinon F, Burns K, McDermott MF, Hawkins PN, Tschopp J: NALP3 forms an IL-1 β -processing inflammasome with increased activity in Muckle-Wells autoinflammatory disorder. *Immunity* 2004;20:319–325.
- Roberts RL, Topless RK, Phipps-Green AJ, Garry RB, Barclay ML, Merriman TR: Evidence of interaction of CARD8 rs2043211 with NALP3 rs35829419 in Crohn's disease. *Genes Immun* 2010;11:351–356.
- Schultz I, Verma D, Halfvarsson J, et al: Combined polymorphisms in genes encoding the inflammasome components NALP3 and CARD8 confer susceptibility to Crohn's disease in Swedish men. *Am J Gastroenterol* 2009;104:1180–1188.
- Kastbom A, Verma D, Eriksson P, Skogh T, Wingren G, Soderkvist P: Genetic variation in proteins of the cryopyrin inflammasome influences susceptibility and severity of rheumatoid arthritis (the Swedish TIRA project). *Rheumatology (Oxford)* 2008;47:415–417.
- Hoffman HM, Mueller JL, Broide DH, Wanderer AA, Kolodner RD: Mutation of a new gene encoding a putative pyrin-like protein causes familial cold autoinflammatory syndrome and Muckle-Wells syndrome. *Nat Genet* 2001;29:301–305.
- Feldmann J, Prieur AM, Quartier P, et al: Chronic infantile neurological cutaneous and articular syndrome is caused by mutations in CIAS1, a gene highly expressed in polymorphonuclear cells and chondrocytes. *Am J Hum Genet* 2002;71:198–203.
- Martinon F, Mayor A, Tschopp J: The inflammasomes: guardians of the body. *Annu Rev Immunol* 2009;27:229–265.
- Tschopp J, Schroder K: NLRP3 inflammasome activation: The convergence of multiple signalling pathways on ROS production? *Nat Rev Immunol* 2010;10:210–215.
- Kanneganti TD, Body-Malapel M, Amer A, et al: Critical role for cryopyrin/NALP3 in activation of caspase-1 in response to viral infection and double-stranded RNA. *J Biol Chem* 2006;281:36560–36568.
- Allen IC, Scull MA, Moore CB, et al: The NLRP3 inflammasome mediates in vivo innate immunity to influenza A virus through recognition of viral RNA. *Immunity* 2009;30:556–565.
- Muruve DA, Petrilli V, Zaiss AK, et al: The inflammasome recognizes cytosolic microbial and host DNA and triggers an innate immune response. *Nature* 2008;452:103–107.
- Munoz-Planillo R, Franchi L, Miller LS, Nunez G: A critical role for hemolysins and bacterial lipoproteins in *Staphylococcus aureus*-induced activation of the NLRP3 inflammasome. *J Immunol* 2009;183:3942–3948.
- Warren SE, Mao DP, Rodriguez AE, Miao EA, Aderem A: Multiple Nod-like receptors activate caspase-1 during *Listeria monocytogenes* infection. *J Immunol* 2008;180:7558–7564.
- Lamkanfi M, Malireddi RK, Kanneganti TD: Fungal zymosan and mannan activate the cryopyrin inflammasome. *J Biol Chem* 2009;284:20574–20581.
- Joly S, Ma N, Sadler JJ, Soll DR, Cassel SL, Sutterwala FS: Cutting edge: *Candida albicans* hyphae formation triggers activation of the NLRP3 inflammasome. *J Immunol* 2009;183:3578–3581.

- 23 Said-Sadier N, Padilla E, Langsley G, Ojcius DM: *Aspergillus fumigatus* stimulates the NLRP3 inflammasome through a pathway requiring ROS production and the Syk tyrosine kinase. *PLoS ONE* 2010;5:e10008.
- 24 Kanneganti TD, Ozoren N, Body-Malapel M, et al: Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/NALP3. *Nature* 2006;440:233–236.
- 25 Gurcel L, Abrami L, Girardin S, Tschopp J, van der Goot FG: Caspase-1 activation of lipid metabolic pathways in response to bacterial pore-forming toxins promotes cell survival. *Cell* 2006;126:1135–1145.
- 26 Pelegrin P, Surprenant A: Pannexin-1 couples to maitotoxin- and nigericin-induced interleukin-1 β release through a dye uptake-independent pathway. *J Biol Chem* 2007;282:2386–2394.
- 27 Dostert C, Guarda G, Romero JF, et al: Malarial hemozoin is an NALP3 inflammasome activating danger signal. *PLoS ONE* 2009;4:e6510.
- 28 Yamasaki K, Muto J, Taylor KR, et al: NLRP3/cryopyrin is necessary for interleukin-1 β (IL-1 β) release in response to hyaluronan, an endogenous trigger of inflammation in response to injury. *J Biol Chem* 2009;284:12762–12771.
- 29 Zhou R, Tardivel A, Thorens B, Choi I, Tschopp J: Thioredoxin-interacting protein links oxidative stress to inflammasome activation. *Nat Immunol* 2010;11:136–140.
- 30 Halle A, Hornung V, Petzold GC, et al: The NALP3 inflammasome is involved in the innate immune response to amyloid- β . *Nat Immunol* 2008;9:857–865.
- 31 Hornung V, Bauernfeind F, Halle A, et al: Silica crystals and aluminum salts activate the NALP3 inflammasome through phagosomal destabilization. *Nat Immunol* 2008;9:847–856.
- 32 Cassel SL, Eisenbarth SC, Iyer SS, et al: The NALP3 inflammasome is essential for the development of silicosis. *Proc Natl Acad Sci USA* 2008;105:9035–9040.
- 33 Duewell P, Kono H, Rayner KJ, et al: NLRP3 inflammasomes are required for atherogenesis and activated by cholesterol crystals. *Nature* 2010;464:1357–1361.
- 34 Thompson NP, Driscoll R, Pounder RE, Wakefield AJ: Genetics versus environment in inflammatory bowel disease: results of a British twin study. *BMJ* 1996;312:95–96.
- 35 Tysk C, Lindberg E, Jarnerot G, Floderus-Myrhed B: Ulcerative colitis and Crohn's disease in an unselected population of monozygotic and dizygotic twins. A study of heritability and the influence of smoking. *Gut* 1988;29:990–996.
- 36 Orholm M, Binder V, Sorensen TI, Rasmussen LP, Kyvik KO: Concordance of inflammatory bowel disease among Danish twins. Results of a nationwide study. *Scand J Gastroenterol* 2000;35:1075–1081.
- 37 Jess T, Riis L, Jespersgaard C, et al: Disease concordance, zygosity, and NOD2/CARD15 status: follow-up of a population-based cohort of Danish twins with inflammatory bowel disease. *Am J Gastroenterol* 2005;100:2486–2492.
- 38 Hanauer SB: Inflammatory bowel disease: epidemiology, pathogenesis, and therapeutic opportunities. *Inflamm Bowel Dis* 2006;12(suppl 1):S3–S9.
- 39 Loftus EV Jr: Clinical epidemiology of inflammatory bowel disease: Incidence, prevalence, and environmental influences. *Gastroenterology* 2004;126:1504–1517.
- 40 European Parliament C: On colours for use in foodstuffs. 1994:13–29.
- 41 Baan R, Straif K, Grosse Y, Secretan B, El Ghissassi F, Coglian V: Carcinogenicity of carbon black, titanium dioxide, and talc. *Lancet Oncol* 2006;7:295–296.
- 42 Lomer MC, Thompson RP, Powell JJ: Fine and ultrafine particles of the diet: influence on the mucosal immune response and association with Crohn's disease. *Proc Nutr Soc* 2002;61:123–130.
- 43 European Food Safety Authority: Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids and Materials in Contact with Food on a request from the commission related to the safety-in-use of rutile titanium dioxide as an alternative to the presently permitted anatase form. *EFSA J* 2004;163:1–12.
- 44 Lomer MC, Hutchinson C, Volkert S, et al: Dietary sources of inorganic microparticles and their intake in healthy subjects and patients with Crohn's disease. *Br J Nutr* 2004;92:947–955.
- 45 Powell JJ, Thoree V, Pele LC: Dietary microparticles and their impact on tolerance and immune responsiveness of the gastrointestinal tract. *Br J Nutr* 2007;98(suppl 1):S59–S63.
- 46 Shepherd NA, Crocker PR, Smith AP, Levi-Strauss DA: Exogenous pigment in Peyer's patches. *Hum Pathol* 1987;18:50–54.
- 47 Powell JJ, Ainley CC, Harvey RS, et al: Characterisation of inorganic microparticles in pigment cells of human gut-associated lymphoid tissue. *Gut* 1996;38:390–395.
- 48 Urbanski SJ, Arsenaault AL, Green FH, Haber G: Pigment resembling atmospheric dust in Peyer's patches. *Mod Pathol* 1989;2:222–226.
- 49 Lomer MC, Thompson RP, Powell JJ: Fine and ultrafine particles of the diet: influence on the mucosal immune response and association with Crohn's disease. *Proc Nutr Soc* 2002;61:123–130.
- 50 Jani PU, McCarthy DE, Florence AT: Titanium dioxide (rutile) particle uptake from the rat GI tract and translocation to systemic organs after oral administration. *Int J Pharm* 1994;105:157–168.
- 51 Ramsden CS, Smith TJ, Shaw BJ, Handy RD: Dietary exposure to titanium dioxide nanoparticles in rainbow trout, (*Oncorhynchus mykiss*): no effect on growth, but subtle biochemical disturbances in the brain. *Eco-toxicology* 2009;18:939–951.
- 52 Bockmann J, Lahl H, Eckert T, Unterhalt B: Blood titanium levels before and after oral administration titanium dioxide. *Pharmazie* 2000;55:140–143.
- 53 Best WR, Beckett JM, Singleton JW, Kern F Jr: Development of a Crohn's disease activity index. National Cooperative Crohn's Disease Study. *Gastroenterology* 1976;70:439–444.
- 54 Lomer MC, Grainger SL, Ede R, et al: Lack of efficacy of a reduced microparticle diet in a multi-centred trial of patients with active Crohn's disease. *Eur J Gastroenterol Hepatol* 2005;17:377–384.
- 55 Winter M, Beer HD, Hornung V, Kramer U, Schins RP, Forster I: Activation of the inflammasome by amorphous silica and TiO(2) nanoparticles in murine dendritic cells. *Nanotoxicology* 2011;5:326–340.
- 56 Yazdi AS, Guarda G, Riteau N, et al: Nanoparticles activate the NLR pyrin domain containing 3 (NLRP3) inflammasome and cause pulmonary inflammation through release of IL-1 α and IL-1 β . *Proc Natl Acad Sci USA* 2010;107:19449–19454.
- 57 Franchi L, Eigenbrod T, Nunez G: Cutting edge: TNF- α mediates sensitization to ATP and silica via the NLRP3 inflammasome in the absence of microbial stimulation. *J Immunol* 2009;183:792–796.
- 58 Chen GY, Nunez G: Sterile inflammation: sensing and reacting to damage. *Nat Rev Immunol* 2010;10:826–837.
- 59 Petrilli V, Papin S, Dostert C, Mayor A, Martinon F, Tschopp J: Activation of the NALP3 inflammasome is triggered by low intracellular potassium concentration. Cell death and differentiation 2007;14:1583–1589.
- 60 Windelborn JA, Lipton P: Lysosomal release of cathepsins causes ischemic damage in the rat hippocampal slice and depends on NMDA-mediated calcium influx, arachidonic acid metabolism, and free radical production. *J Neurochem* 2008;106:56–69.
- 61 Li Z, Berk M, McIntyre TM, Gores GJ, Feldstein AE: The lysosomal-mitochondrial axis in free fatty acid-induced hepatic lipotoxicity. *Hepatology* 2008;47:1495–1503.
- 62 Monteleone G, Trapasso F, Parrello T, et al: Bioactive IL-18 expression is up-regulated in Crohn's disease. *J Immunol* 1999;163:143–147.
- 63 Pizarro TT, Michie MH, Bentz M, et al: IL-18, a novel immunoregulatory cytokine, is up-regulated in Crohn's disease: expression and localization in intestinal mucosal cells. *J Immunol* 1999;162:6829–6835.
- 64 Moyes SM, Smyth SH, Shipman A, Long S, Morris JF, Carr KE: Parameters influencing intestinal epithelial permeability and microparticle uptake in vitro. *Int J Pharm* 2007;337:133–141.
- 65 Koenenman BA, Zhang Y, Westerhoff P, Chen Y, Crittenden JC, Capco DG: Toxicity and cellular responses of intestinal cells exposed to titanium dioxide. *Cell Biol Toxicol*;26: 225–238.
- 66 Tschopp J, Martinon F, Burns K: NALPs: a novel protein family involved in inflammation. *Nat Rev Mol Cell Biol* 2003;4:95–104.